

For Research Use Only

Mouse CD20 Polyclonal antibody, PBS Only

Catalog Number:31909-1-PBS



Basic Information

Catalog Number:

31909-1-PBS

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG36918

GenBank Accession Number:

NM_007641

GeneID (NCBI):

12482

UNIPROT ID:

P19437

Full Name:

membrane-spanning 4-domains, subfamily A, member 1

Calculated MW:

32 kDa

Observed MW:

35 kDa

Purification Method:

Antigen affinity Purification

Applications

Tested Applications:

WB, IHC, IF-P, Indirect ELISA

Species Specificity:

mouse, rat

Background Information

CD20 is a 33-37 kDa transmembrane phosphoprotein. CD20 is a B-lymphocyte surface molecule that is widely expressed during B-cell ontogeny, from early pre-B-cell developmental stages until final differentiation into plasma cells. CD20 functions as a calcium-permeable cation channel. It is involved in the regulation of B-cell activation and proliferation. CD20 serves as a useful target for antibody-mediated therapeutic depletion of B-cells.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 31909-1-AP (CD20 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 31909-1-AP (CD20 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using 31909-1-AP (CD20 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse colon tissue using CD20 antibody (31909-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using CD20 antibody (31909-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat spleen tissue using CD20 antibody (31909-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 31909-1-PBS in a different storage buffer formulation.