For Research Use Only

LSM2 Polyclonal antibody Catalog Number: 32074-1-AP




Basic Information	Catalog Number: 32074-1-AP	GenBank Accession Number: BC009192	Purification Method: Antigen affinity Purification	
	Size: 750 ug/ml	GenelD (NCBI): 57819	Recommended Dilutions: WB 1:2000-1:12000	
	Source: Rabbit	UNIPROT ID: Q9Y333		
	Isotype: IgG Immunogen Catalog Number:	Full Name: LSM2 homolog, U6 small nuclear RNA associated (S. cerevisiae) Calculated MW: 11 kDa		
	AG34865			
		Observed MW: 11 kDa		
Applications	Tested Applications: WB, ELISA	Positive Controls:		
	Species Specificity: human	WB : HEK-	293 cells, K-562 cells, human placenta tissue	
Background Information	Sm-like proteins contain the Sm s that folds as a loop. The Sm-like p which are important for pre-mRNA stabilizes the 3' end of the splic snRNA in vitro and interact with th	Sm-like proteins were identified in a variety of organisms based on sequence homology with the Sm protein family. Sm-like proteins contain the Sm sequence motif, which consists of 2 regions separated by a linker of variable length that folds as a loop. The Sm-like proteins are thought to form a stable heteromer present in tri-snRNP particles, which are important for pre-mRNA splicing (PMID:15075370). Lsm2-Lsm8 (Lsm stands for Like Sm), binds and stabilizes the 3' end of the spliceosomal U6 snRNA Lsm proteins facilitate the annealing of U6 snRNA with U4 snRNA in vitro and interact with the U4/U6 annealing factor Prp24, suggesting that Lsm proteins function in U4/U6 snRNP formation in vivo (PMID:15485930, 17251193).		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptg W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32074-1-AP (LSM2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.