For Research Use Only

CD3 Delta Polyclonal antibody

Catalog Number:32144-1-AP




Basic Information	Catalog Number: 32144-1-AP	GenBank Accession Number: BC070321	Purification Method: Antigen affinity Purification		
	Size: 450 ug/ml Source: Rabbit Isotype: IgG	GeneID (NCBI): 915 UNIPROT ID: P04234	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:400-1:1600		
				Full Name: CD3d molecule, delta (CD3-TCR complex) Calculated MW: 171 aa, 19 kDa	
		Observed MW: 21 kDa			
		Applications	Tested Applications:	Positive Controls: WB : Jurkat cells, MOLT-4 cells IHC : human tonsillitis tissue,	
			WB, IHC, ELISA Species Specificity: human		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T- lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells.				
Storage	Storage: Store at -20°C. Stable for on Storage Buffer: PBS with 0.02% sodium azio Aliquoting is unnecessary fo	de and 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 32144-1-AP (CD3 Delta antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 32144-1-AP (CD3 Delta antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.