

For Research Use Only

VE-cadherin/CD144 Polyclonal antibody

Catalog Number: 32332-1-AP

1 Publications



Basic Information

Catalog Number: 32332-1-AP	GenBank Accession Number: NM_009868.4	Purification Method: Antigen affinity Purification
Source: Rabbit	GeneID (NCBI): 12562	Recommended Dilutions: WB: 1:2000-1:10000 IF/ICC: 1:50-1:500
Isotype: IgG	UNIPROT ID: P55284	
Immunogen Catalog Number: EG0921	Full Name: cadherin 5	
	Calculated MW: 88kDa	
	Observed MW: 130 kDa, 100 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls: WB : bEnd.3 cells, mouse lung tissue, rat lung tissue IF/ICC : bEnd.3 cells,
Cited Applications: WB, IF	
Species Specificity: mouse, rat	
Cited Species: mouse	

Background Information

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. Vascular endothelial cadherin (VE-cadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of the type II classical cadherin family of cell adhesion proteins (PMID: 21269602). VE-cadherin is expressed specifically in endothelial cells and mediates homophilic adhesion in the vascular endothelium (PMID: 1522121; 8555485; 21269602). VE-cadherin plays a role in the organization of lateral endothelial junctions and in the control of permeability properties of vascular endothelium (PMID: 1522121). VE-cadherin has also been shown to be required for angiogenesis (PMID: 16473763; 18162609). The calculated molecular weight of VE-cadherin is 88 kDa and the apparent molecular weight of 120-140 kDa is higher due to post-translational glycosylation and phosphorylation (PMID: 10460833; 29894844). Full-length VE-cadherin can be proteolytically cleaved to generate a fragment of 90-100 kDa (PMID: 9786462; 22064597).

Notable Publications

Author	Pubmed ID	Journal	Application
Minghao Luo	41240537	Phytomedicine	WB, IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32332-1-AP (VE-cadherin/CD144 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed bEnd.3 cells using VE-cadherin/CD144 antibody (32332-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).