For Research Use Only

CD13 Polyclonal antibody

Catalog Number: 32752-1-AP



Basic Information

Catalog Number:

32752-1-AP

NM_008486.2

Concentration:

450 ug/ml

Source:

Rabbit

P97449

Isotype:

GenBank Accession Number:

NM_008486.2

GeneID (NCBI):

16790

UNIPROT ID:

P97449

Full Name:

alanyl (membrane) aminopeptidase

Immunogen Catalog Number:Calculated MW:EG3592110 kDaObserved MW:

150 kDa

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Antigen affinity Purification Recommended Dilutions: WB 1:2000-1:10000 IHC 1:200-1:800

Purification Method:

Positive Controls:

WB: mouse kidney tissue, mouse liver tissue, mouse

small intestine tissue

 $\label{eq:IHC:mouse} \textbf{IHC:} mouse spleen tissue,$

Background Information

CD13, also named as APN, ANPEP(aminopeptidase N) or PEPN, is belongs to the peptidase M1 family. CD13 is a heavily glycosylated, ~150-240 kDa, type-II membrane, expressed by most cells of myeloid origin including monocytes, macrophages, granulocytes, and their hematopoietic precursors. It is also abundantly expressed in the brush border of epithelial cells from renal proximal tubules and small intestine, in prostatic epithelial cells, in bile duct canaliculi, in mast cells, and, in some cases, in fibroblasts and smooth muscle cells. CD13 is a multifunctional protein and plays varying roles in cell migration, cell proliferation, cell differentiation and so on. CD13 participates in angiogenesis generating and modulating angiogenic signals, and can be a marker of angiogenic vessels. CD13 is also a pan-myeloid marker, present on mature granulocytes and monocytes. (PMID: 8805662, 10098327, 18603472, 18097955, 17897790, 17888402, 21339174)

Storage


Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 32752-1-AP (CD13 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 32752-1-AP (CD13 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 32752-1-AP (CD13 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).