For Research Use Only

GALC Polyclonal antibody

Catalog Number:51051-2-AP 3 Publications




Basic Information	Catalog Number: 51051-2-AP	GenBank Accession Numbe BC036518	r: Purification Method: Antigen affinity purification
	Size:	GeneID (NCBI):	Recommended Dilutions:
	400 µg/ml	2581	WB 1:500-1:2000
	Source:	UNIPROT ID:	IHC 1:20-1:200
	Rabbit	P54803	IF/ICC 1:50-1:500
	Isotype:	Full Name:	
	IgG	galactosylceramidase	
	Immunogen Catalog Number: Calculated MW: AG0483 668 aa, 76 kDa		
		Observed MW: 30 kDa	
		50 100	
Applications	Tested Applications:	Posi	tive Controls:
	IF/ICC, IHC, WB,ELISA	WB	: human brain tissue,
	Cited Applications: WB	IHC	: human gliomas tissue,
		IF/I	CC : SH-SY5Y cells,
	Species Specificity: human, mouse		
	Cited Species:		
	human		
	Note-IHC: suggested ar with TE buffer pH 9.0; (* antigen retrieval may b) Alternatively, e performed	
	with citrate buffer pH 6.	•	
Background Information	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an	lysosomal enzyme involved in a lelial cells of the small intestine l into 50 and 30 kDa subunits. Mu autosomal recessive neurodege	itation of GALC gene will lead to Krabbe disease
	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an	lysosomal enzyme involved in a lelial cells of the small intestine l into 50 and 30 kDa subunits. Mu autosomal recessive neurodege	and colon. The normal GALC mRNA encodes the station of GALC gene will lead to Krabbe disease enerative disorder that affects both the central ar
	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an peripheral nervous system due to	lysosomal enzyme involved in felial cells of the small intestine linto 50 and 30 kDa subunits. Mu autosomal recessive neurodege an enzymatic defect of galactoo Pubmed ID Journal	and colon. The normal GALC mRNA encodes the station of GALC gene will lead to Krabbe disease enerative disorder that affects both the central ar cerebrosidase. This antibody is specific to GALC. Application
	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an peripheral nervous system due to Author	lysosomal enzyme involved in telial cells of the small intestine linto 50 and 30 kDa subunits. Mu autosomal recessive neurodege an enzymatic defect of galactoo Pubmed ID Journal	and colon. The normal GALC mRNA encodes the station of GALC gene will lead to Krabbe disease enerative disorder that affects both the central ar cerebrosidase. This antibody is specific to GALC. Application wet WB
	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an peripheral nervous system due to Author Mohammad Arif Hossain	lysosomal enzyme involved in telial cells of the small intestine linto 50 and 30 kDa subunits. Mu autosomal recessive neurodege an enzymatic defect of galactoo Pubmed ID Journal 26108143 J Hum Gen	and colon. The normal GALC mRNA encodes the station of GALC gene will lead to Krabbe disease enerative disorder that affects both the central and cerebrosidase. This antibody is specific to GALC. Application wet WB
Notable Publications	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an peripheral nervous system due to Author Mohammad Arif Hossain Jagoda Litowczenko Hossain Mohammad Arif MA	lysosomal enzyme involved in telial cells of the small intestine linto 50 and 30 kDa subunits. Mu autosomal recessive neurodege an enzymatic defect of galactoo Pubmed ID Journal 26108143 J Hum Gen 37862812 Biomater A	and colon. The normal GALC mRNA encodes the itation of GALC gene will lead to Krabbe disease enerative disorder that affects both the central ar cerebrosidase. This antibody is specific to GALC. Application et WB
Background Information Notable Publications	Galactosylceramidase (Galc) is a lipid in myelin, kidney, and epith kDa precursor, which is processed (globoid cell leukodystrophy), an peripheral nervous system due to Author Mohammad Arif Hossain Jagoda Litowczenko	lysosomal enzyme involved in telial cells of the small intestine linto 50 and 30 kDa subunits. Mu autosomal recessive neurodege an enzymatic defect of galactoo Pubmed ID Journal 26108143 J Hum Gen 37862812 Biomater A 24252386 Gene	and colon. The normal GALC mRNA encodes the itation of GALC gene will lead to Krabbe disease enerative disorder that affects both the central ar cerebrosidase. This antibody is specific to GALC. Application et WB

 For technical support and original validation data for this product please contact:


 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 51051-2-AP (GALC antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas using 51051-2-AP (GALC antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using 51051-2-AP (CALC antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).