## For Research Use Only

# IkB Alpha Polyclonal antibody

Catalog Number:51066-1-AP 17 Publications



**Basic Information** 

Catalog Number: 51066-1-AP

Size: 300 μg/ml Source: Rabbit

Isotype:

IgG

Calculated MW: 36 kDa Observed MW: 39 kDa

inhibitor, alpha

BC002601

4792

P25963

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

GenBank Accession Number:

nuclear factor of kappa light

polypeptide gene enhancer in B-cells

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

WB 1:500-1:2000

**Applications** 

**Tested Applications:** WB, IP, IHC, ELISA

Cited Applications:

Species Specificity: human, mouse **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: PC-3 cells, HeLa cells, Jurkat cells

IP: HeLa cells.

IHC: human cervical cancer tissue, human liver tissue

## **Background Information**

Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (NFKBIA, synonyms: IKBA, MAD-3, NFKB1). NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by  $I-kappa-B\ proteins\ (NFKBIA\ or\ NFKBIB),\ which\ inactivate\ NF-kappa-B\ by\ trapping\ it\ in\ the\ cytoplasm.$ Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5prime HGGARNYYCC 3-prime.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Zheng Zhenhua Z	21810613	J Immunol	WB
Hao Wang	34745126	Front Immunol	WB
Mathilde Chataigner	33801489	Nutrients	WB

Storage

Storage: Store at -20°C. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


#### **Selected Validation Data**




PC-3 cells were subjected to SDS PAGE followed by western blot with 51066-1-AP (lkB alpha antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-IkB Alpha (IP:51066-1-AP, 4ug; Detection:51066-1-AP 1:1000) with HeLa cells lysate 3200ug.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 51066-1-AP (IkB alpha antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 51066-1-AP (IkB alpha antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).