For Research Use Only

USP22 Polyclonal antibody

Catalog Number: 55110-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number: GenBank Accession Number: 55110-1-AP NM_015276

Concentration: GeneID (NCBI): 23326

Source: UNIPROT ID: Rabbit Q9UPT9

Isotype: Full Name:

ubiquitin specific peptidase 22

Calculated MW: 60 kDa Observed MW: 58-60 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications: WB, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP

Species Specificity: human

Cited Species: human, mouse, rat Positive Controls:

WB: A549 cells, HepG2 cells

IP: HepG2 cells,

Background Information

USP22, also named as KIAA1063 and USP3L, belongs to the peptidase C19 family and UBP8 subfamily. It is histone deubiquitinating component of the transcription regulatory histone acetylation (HAT) complex SAGA. USP22 catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. It is recruited to specific gene promoters by activators such as MYC, where it is required for transcription. USP22 is required for nuclear receptor-mediated transactivation and cell cycle progression. USP22 has 2 isoforms with molecular mass of 58kDa and 60 kDa. The antibody is specific to USP22.

Notable Publications

Author	Pubmed ID	Journal	Application
Dianyun Ren	34743406	Mol Oncol	WB,IP
Zelong Han	36333288	Cell Death Discov	WB,IHC,IF,IP
Qingxiang Liu	29732405	Sci Adv	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 55110-1-AP (USP22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 55110-1-AP (USP22 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 55110-1-AP (USP22 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of USP22 antibody (55110-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP22 transfected A549 cells.



IP result of anti-USP22 (IP:55110-1-AP, 4ug; Detection:55110-1-AP 1:500) with HepG2 cells lysate 4000 ug.