

For Research Use Only

EGR1 Polyclonal antibody

Catalog Number: 55117-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number:

55117-1-AP

Size:

450 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001964

GeneID (NCBI):

1958

UNIPROT ID:

P18146

Full Name:

early growth response 1

Calculated MW:

58 kDa

Observed MW:

70-80 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A549 cells, MCF-7 cells, PC-3 cells

IHC : human placenta tissue, human cervical cancer tissue

IF/ICC : MCF-7 cells,

Background Information

Early growth response 1 (EGR1) is a member of EGR family which are transcriptional factors that contain three repetitive zinc finger DNA binding domains which bind to EGR response elements (ER) to regulate target gene expression. The expression of EGR family members is induced by growth factors, with EGR1 expression being induced by NGF. Increased EGR1 expression activates transcription of other signaling molecules, including CDK5 and tyrosine hydroxylase, and exerts long term effects on neural cell growth and differentiation. Egr-1 binds to the DNA sequence 5'-CGCCCCGC-3' (Egr-site), thereby activating transcription of target genes whose products are required for mitogenesis and differentiation. Western blotting of nuclear and cytoplasmic fractions of CNS tissue verified the cytoplasmic Egr1 as 110-kDa dimer or after denaturation as a 57-kDa version, which is reflecting its theoretical molecular weight of 57 kDa. In the nucleus Egr1 is found as 75-kDa homo- or heterodimer and as 35-kDa variant under strong denaturation.

Notable Publications

Author	Pubmed ID	Journal	Application
Shuyue Sheng	34608568	In Vitro Cell Dev Biol Anim	WB
Yan-Lin Yang	30462533	Am J Physiol Cell Physiol	WB
Yin Wang	32444791	Nat Commun	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 55117-1-AP (EGR1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 55117-1-AP (EGR1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human EGR1 (55117-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using EGR1 antibody (55117-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using EGR1 antibody (55117-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).