

CD30/TNFRSF8 Polyclonal antibody

Catalog Number: 55132-1-AP

Basic Information

Catalog Number:

55132-1-AP

Size:

200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001243

GeneID (NCBI):

943

UNIPROT ID:

P28908

Full Name:tumor necrosis factor receptor
superfamily, member 8**Calculated MW:**

64 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IHC 1:500-1:2000

IF-P 1:50-1:500

Applications

Tested Applications:

IHC, IF-P, ELISA

Species Specificity:

human

Positive Controls:

IHC : human lymphoma tissue, human tonsillitis tissue

IF-P : human lymphoma tissue,

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Background Information

CD30, also known as TNFRSF8 or Ki-1, is a transmembrane glycoprotein belonging to the tumor necrosis factor receptor superfamily. It is composed of an extracellular domain with six cysteine-rich repeats, a transmembrane segment, and an intracellular domain (PMID: 1310894). CD30 is expressed on a small subset of activated T and B lymphocytes, and a variety of lymphoid neoplasms, with the highest expression in classical Hodgkin's lymphoma and anaplastic large cell lymphoma (PMID: 9826579; 28885612). The CD30 ligand (CD30L or CD153) is a type II transmembrane glycoprotein of the tumor necrosis factor ligand superfamily (PMID: 8701986). Signal transduction of CD30/CD30L utilizes signal transducers, TNFR-associated factors (TRAF1, 2, 3 and 5) (PMID: 9826579).

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:0.1M NaHCO₃, 0.1M glycine, 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human T-cell anaplastic large cell lymphoma (ALCL) tissue slide using 55132-1-AP (CD30 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human T-cell anaplastic large cell lymphoma (ALCL) tissue slide using 55132-1-AP (CD30 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human T-cell anaplastic large cell lymphoma (ALCL) tissue using CD30 antibody (55132-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human T-cell anaplastic large cell lymphoma (ALCL) tissue using CD30 antibody (55132-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).