For Research Use Only

PTBP1 Polyclonal antibody

Catalog Number: 55181-1-AP

Featured Product

1 Publications



Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

IF-P 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 55181-1-AP Size:

Size: 350 µg/ml Source: Rabbit Isotype: GenBank Accession Number:

NM_002819 GeneID (NCBI): 5725 UNIPROT ID: P26599 Full Name:

Full Name: polypyrimidine tract binding protein

1 Calculated MW:

60 kDa Observed MW: 55-62 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications:

WB

Species Specificity: human, mouse Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, HeLa cells
IHC: mouse brain tissue,
IF-P: mouse brain tissue,
IF/ICC: HEK-293 cells,

Background Information

Polypyrimidine tract-binding protein (PTBP1), also known as heterogeneous nuclear ribonucleoprotein type I (hnRNP I), is a nuclear protein that binds pre-mRNAs in specific regions of the hnRNA-protein complexes sensitive to micrococcal nuclease. It involves in pre-mRNA splicing and the regulation of alternative splicing events. PTBP1 promotes RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA, and promotes the binding of U2 snRNP to pre-mRNA. Besides, it can cooperate with RAVER1 to modulate switching between mutually exclusive exons during maturation of the TPM1 pre-mRNA, and represses the splicing of MAPT/Tau exon 10

Notable Publications

Author	Pubmed ID	Journal	Application
Shaojun Zhang	34737357	Cell Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 55181-1-AP (PTBP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




HepG2 cells were subjected to SDS PAGE followed by western blot with 55181-1-AP (PTBP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of PTBP1 antibody (55181-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PTBP1 transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 55181-1-AP (PTBP1 antibody), at dilution of 1:100 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of HEK-293 cells using 55181-1-AP (PTBP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).