For Research Use Only

HLA-F Polyclonal antibody

Catalog Number: 55379-1-AP



Basic Information

Catalog Number: 55379-1-AP

 $\begin{array}{lll} \text{55379-1-AP} & \text{NM_018950} \\ \text{Size:} & \text{GeneID (NCBI):} \\ 1000 \ \mu \ \text{g/ml} & 3134 \\ \text{Source:} & \text{UNIPROT ID:} \\ \text{Rabbit} & \text{P30511} \\ \end{array}$

Isotype: Full Name:

G major histocompatibility complex,

class I, F
Calculated MW:

50 kDa

Observed MW:
50 kDa, 66-70 kDa

GenBank Accession Number:

Applications

Tested Applications: WB,ELISA

Species Specificity: human, mouse

Positive Controls:

WB: Raji cells, A431 cells, human placenta tissue,

Purification Method:

WB 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

Jurkat cells, mouse spleen tissue

Background Information

Human major histocompatibility complex (MHC) antigens, also referred to as human leukocyte antigens (HLA), are encoded by genes located on the short arm of chromosome 6 (6p21.3). There are two classes of HLA antigens: class I and class II. This class I molecules are membrane glycoproteins composed of a heavy (alpha) chain which is encoded by a HLA class I gene, and β 2-microglobulin light (beta) chain. The most extensively characterized members of the HLA class I gene family are the genes encoding the major transplantation antigenes, HLA-A, B and C. HLA-F is a non-classical MHC class I molecule. (PMID: 667938; 3375250; 2249951)


Storage

Storage: Store at -20°C. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Raji cells were subjected to SDS PAGE followed by western blot with 55379-1-AP (HLA-F antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.