## For Research Use Only

# **HLA-A Polyclonal antibody**

Catalog Number: 55383-1-AP 2 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 55383-1-AP NM\_002116 GeneID (NCBI): Concentration: 500 ug/ml 3105 **UNIPROT ID:** Source: Rabbit P04439 Full Name: Isotype: major histocompatibility complex,

IF/ICC 1:50-1:500

Calculated MW: 41 kDa Observed MW: 41-44 kDa

class I. A

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications:

IHC, IF

Species Specificity: human

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells, A431 cells

IHC: human tonsillitis tissue, human small intestine

**Purification Method:** 

WB 1:1000-1:6000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

tissue

IF/ICC: HepG2 cells,

# **Background Information**

Human major histocompatibility complex (MHC) antigens, also referred to as human leukocyte antigens (HLA), are encoded by genes located on the short arm of chromosome 6 (6p21.3). There are two classes of HLA antigens: class I (HLA-A, B and C) and class II (HLA-D). This class I molecules are polymorphic membrane glycoproteins composed of a heavy (alpha) chain (44 kDa) which is encoded by a HLA class I gene (HLA-A, B or C), and β 2-microglobulin light (beta) chain (12 kDa). They are involved in the presentation of foreign antigens to the immune system. (PMID: 667938; 3375250)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lucas A Tavares	31915283	J Virol	IF
Yanling Wang	38694916	Front Pharmacol	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 55383-1-AP (HLA-A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 55383-1-AP (HLA-A antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 55383-1-AP (HLA-A antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using HLA-A antibody (55383-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).