For Research Use Only

NR4A3 Polyclonal antibody, PBS Only

Catalog Number:55405-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

55405-1-PBS

NM_006981 GeneID (NCBI): Antigen affinity purification

Size: 1 mg/ml Source:

8013 **UNIPROT ID:** Q92570

Rabbit Isotype:

nuclear receptor subfamily 4, group A,

member 3 Calculated MW:

Full Name:

70 kDa Observed MW: 68 kDa

Applications

Tested Applications: WB, IF/ICC, Indirect ELISA Species Specificity:

human, mouse

Background Information

Nuclear receptor 4A3 (NR4A3, also known as NOR-1), NR4A1 and NR4A2 are TFs belonging to NR4A orphan nuclear $receptor \, subfamily. \, The \, role \, of \, all \, three \, NR4A \, members \, in \, regulating \, gene \, expression \, involved \, in \, cell \, proliferation,$ differentiation, apoptosis, inflammation and metabolism in cardiovascular diseases has been widely studied. In osteoblasts, NR4A receptor has been proved to regulate the expression and differentiation of osteoblast genes. It is reported that NR4A receptor is highly expressed in human coronary atherosclerotic lesions, and NR4A3 deficiency can reduce the proliferation and migration of VSMCs and the formation of neointima after vascular injury. Therefore, NR4A3 may play an important role in mediating phenotypic transformation of VSMC (PMID: 38629274). It is expected to be located in nucleus. The protein has three isforms, isoform alpha is highly expressed in skeletal muscle. Isoform beta is highly expressed in skeletal muscle and low expressed in fetal brain and placenta. The calculated molecular weight of NR4A3 is 68 kDa.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


Selected Validation Data



mouse spleen tissue were subjected to SDS PAGE followed by western blot with 55405-1-AP (NR4A3 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 55405-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 55405-1-AP (NR4A3 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 55405-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed ROS1728 cells using NR4A3 antibody (55405-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 55405-1-PBS in a different storage buffer formulation.