For Research Use Only

KHSRP Polyclonal antibody

Catalog Number:55409-1-AP

Featured Product

9 Publications





Basic Information	Catalog Number: 55409-1-AP			Purification Method: Antigen affinity purification	
	Size: Genel D (NCBI): 800 µ g/ml 8570			Recommended Dilutions: WB 1:1000-1:6000	
	Source:UNIPROT ID:RabbitQ92945				
	Isotype:Full Name:IgGKH-type splicing regulatory protein				
	Calculated MW: 73 kDa				
	Observed MW: 75 kDa				
Applications	Tested Applications:		Positive Controls:		
	WB, ELISA Cited Applications: WB, IF, RIP		WB : HEK-293T cells, HeLa cells, HepG2 cells, Jurkat cells, MCF-7 cells, mouse brain tissue, rat brain tissue		
	Species Specificity: human, mouse, rat				
	Cited Species: human, mouse				
Background Information	KHSRP, also named as FUBP2, KSRP, and p75, belongs to the KHSRP family. It binds to the dendritic targeting element and may play a role in mRNA trafficking. KHSRP is a part of a ternary complex that binds to the downstream control sequence (DCS) of the pre-mRNA. It mediates exon inclusion in transcripts that are subject to tissue-specific alternative splicing. KHSRP may interact with single-stranded DNA from the far-upstream element (FUSE). May activate gene expression. It is involved in the degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly by recruiting degradation machinery to ARE-containing mRNAs.				
Notable Publications	Author	Pubmed ID Jo	ournal	Application	
	Huajun Jiang	26523989 A	therosclerosis	WB	
	Xiaomei He	34614161 N	ucleic Acids Res	WB	
	Zhenzhen Chen	34320348 C	ell Rep	WB,RIP	
Storage	Storage: Store at -20°C. Stable for or	ne year after shipment.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 55409-1-AP (KHSRP antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. WB result of KHSRP antibody (55409-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-KHSRP transfected HeLa cells.