## For Research Use Only

# RAB7A Polyclonal antibody

Catalog Number: 55469-1-AP

Featured Product

63 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 55469-1-AP NM\_004637 GeneID (NCBI): Concentration: 1000 ug/ml 7879 **UNIPROT ID:** Source: Rabbit P51149 Full Name: Isotype:

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500 IF/ICC 1:50-1:500

RAB7A, member RAS oncogene family

Calculated MW: 23 kDa Observed MW: 23 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse, rat **Cited Species:** 

human, mouse, rat, pig, monkey, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: L-929 cells, NIH/3T3 cells, A431 cells, SH-SY5Y cells, RAW 264.7 cells, C6 cells, mouse brain tissue, rat brain tissue

IHC: human kidney tissue, human skeletal muscle tissue, human stomach tissue

IF/ICC: HeLa cells,

# **Background Information**

RAB7A, also named as RAB7, belongs to the small GTPase superfamily and Rab family. It is involved in late endocytic transport. RAB7A contributes to the maturation of phagosomes (acidification). Defects in RAB7A are the cause of Charcot-Marie-Tooth disease type 2B (CMT2B). This antibody is specific to RAB7A.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Wenmin Yu	34676302	Open Med (Wars)	IHC
Wenqing Li	31578588	Oncol Rep	
Denghui Wei	32958903	Cell Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 55469-1-AP (RAB7A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 55469-1-AP (RAB7A antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 55469-1-AP (RAB7A antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using RAB7A antibody (55469-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human RAB7A (55469-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).