

For Research Use Only

CD2 Monoclonal antibody

Catalog Number: 60005-1-Ig **1 Publications**



Basic Information

Catalog Number: 60005-1-Ig	GenBank Accession Number: BC033583	Purification Method: Protein G purification
Size: 1800 µg/ml	GeneID (NCBI): 914	CloneNo.: 3A10B2
Source: Mouse	Full Name: CD2 molecule	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Isotype: IgG1	Calculated MW: 351 aa, 39 kDa	
Immunogen Catalog Number: AG0423	Observed MW: 50 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB: Jurkat cells, human spleen tissue IHC: human tonsillitis tissue,
Cited Applications: IF	
Species Specificity: human	
Cited Species: human, sheep	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD2 is a cell surface glycoprotein present on a majority of thymocytes, all mature T cells and subset of NK cells but not on B lymphocytes. It is a pan T-cell marker. CD2 interacts with lymphocyte function-associated antigen (LFA-3/CD58) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Notable Publications

Author	Pubmed ID	Journal	Application
Tomoyuki Abe	33395577	Exp Hematol	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Untreated and PNGase F-treated lysates of Jurkat cells were subjected to SDS PAGE followed by western blot with 60005-1-Ig (CD2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).



Jurkat cells were subjected to SDS PAGE followed by western blot with 60005-1-Ig (CD2 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60005-1-Ig (CD2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60005-1-Ig (CD2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).