For Research Use Only

TGFBI / BIGH3 Monoclonal antibody, PBS Only



Purification Method:

precipitation

CloneNo.: 3E11D11

Caprylic acid/ammonium sulfate

Catalog Number: 60007-1-PBS

Featured Product

Basic Information

Catalog Number:

60007-1-PBS

Size: 1mg/ml

Source: Mouse

Isotype: IgG2a

Immunogen Catalog Number:

AG0241

GenBank Accession Number:

BC000097

GeneID (NCBI): 7045

ENSEMBL Gene ID:

ENSG00000120708

UNIPROT ID:

Q15582

Full Name: transforming growth factor, beta-

induced, 68kDa

Calculated MW: 683 aa, 75 kDa Observed MW: 68 kDa

Applications

Tested Applications:

WB, IP, IF, IHC, Indirect ELISA

Species Specificity:

human

Background Information

TGFBI, also named as BIGH3, Kerato-epithelin and RGD-CAP, binds to type I, II, and IV collagens. TGFBI is an adhesion protein which may play an important role in cell-collagen interactions. In cartilage, it may be involved in endochondral bone formation. TGFBI is an extracellular matrix adaptor protein, it has been reported to be differentially expressed in transformed tissues. TGFBI is a predictive factor of the response to chemotherapy, and suggest the use of TGFBI-derived peptides as possible therapeutic adjuvants for the enhancement of responses to chemotherapy. (PMID:20509890) Defects in TGFBI are the cause of epithelial basement membrane corneal dystrophy (EBMD). Defects in TGFBI are the cause of corneal dystrophy active type 1 (CDL1). Defects in TGFBI are a cause of corneal dystrophy Thiel-Behnke type (CDTB). Defects in TGFBI are the cause of Reis-Buecklers corneal dystrophy (CDRB). Defects in TGFBI are the cause of lattice corneal dystrophy type 3A (CDL3A). Defects in TGFBI are the cause of Avellino corneal dystrophy (ACD).

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only


Selected Validation Data




human kidney tissue were subjected to SDS PAGE followed by western blot with 60007-1-lg (TGFBI / BIGH3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60007-1-Ig (TGFBI / BIGH3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60007-1-1g (TGFBI / BIGH3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.



IP result of anti-TGFBI / BIGH3 (IP:60007-1-Ig, 4ug; Detection:60007-1-Ig 1:300) with HeLa cells lysate 1200ug. This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using TGFBI / BIGH3 antibody (60007-1-Ig, Clone: 3E11D11) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using TGFBI / BIGH3 antibody (60007-1-Ig, Clone: 3E11D11) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60007-1-PBS in a different storage buffer formulation.