For Research Use Only

Beta Actin Monoclonal antibody

Catalog Number:60008-2-lg 9 Publications



Basic Information	Catalog Number: 60008-2-1g	GenBank Accession Number: BC002409	Purification Method: Caprylic Acid/Ammonium Sulfate
	Size:	GenelD (NCBI):	Precipitation
	500 µg/ml	60	CloneNo.:
	Source: Mouse	UNIPROT ID: P60709	7A9H3
	Isotype: IgM	Full Name: actin, beta	
	Immunogen Catalog Number: AG0297	Calculated MW: 375 aa, 42 kDa	
		Observed MW: 45 kDa	
Applications	Tested Applications: ELISA		
	Cited Applications: WB		
	Species Specificity: human		
	Cited Species: human, mouse		
Background Informatior	Beta actin, also named as ACTB and F-Actin, belongs to the actin family. Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. least six isoforms of actins are known in mammals and other vertebrates: alpha (ACTC1, cardiac muscle 1), alpha (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. This antibody can recogni all of actins.		
	enteric smooth muscle). Beta and ACTG2 (rich in muscles) has 375a widely used as the internal contro	a and ACTG1(found in non-muscle	n proteins. Most actins consist of 376aa, while cells) has only 374aa. Beta actin has been
Notable Publications	enteric smooth muscle). Beta and ACTG2 (rich in muscles) has 375a widely used as the internal contro all of actins.	a and ACTG1(found in non-muscle	n proteins. Most actins consist of 376aa, while cells) has only 374aa. Beta actin has been
lotable Publications	enteric smooth muscle). Beta and ACTG2 (rich in muscles) has 375a widely used as the internal contro all of actins. Author I	a and ACTG1(found in non-muscle ol in RT-PCR and Western Blotting	n proteins. Most actins consist of 376aa, while cells) has only 374aa. Beta actin has been as a 42-kDa protein. This antibody can recogni
lotable Publications	enteric smooth muscle). Beta and ACTG2 (rich in muscles) has 375a widely used as the internal contro all of actins. Author I Su Yao	a and ACTG1(found in non-muscle ol in RT-PCR and Western Blotting Pubmed ID Journal	n proteins. Most actins consist of 376aa, while e cells) has only 374aa. Beta actin has been as a 42-kDa protein. This antibody can recogni Application
Notable Publications	enteric smooth muscle). Beta and ACTG2 (rich in muscles) has 375a widely used as the internal contro all of actins. Author I Su Yao 2 Su Yao 2	a and ACTG1(found in non-muscle ol in RT-PCR and Western Blotting Pubmed ID Journal 35359405 Front Oncol	n proteins. Most actins consist of 376aa, while cells) has only 374aa. Beta actin has been as a 42-kDa protein. This antibody can recogni Application

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer. Selected Validation Data