For Research Use Only

GATA1 Monoclonal antibody

Catalog Number:60011-1-lg Featured Product 7 Publications





	Catalog Number: 60011-1-lg	BC009797	on Number:	Purification Method: Protein A purification	
	Size:	GeneID (NCBI):		CloneNo.:	
	1000 ug/ml	2623		5E2A8	
	Source:	UNIPROT ID:		Recommended Dilutions:	
	Mouse	P15976 WB 1:1000-1:8000 Full Name: IHC 1:500-1:2000 GATA binding protein 1 (globin transcription factor 1)			
	Isotype: IgG1				
	Immunogen Catalog Number: AG1350				
		Calculated MW: 43 kDa			
		Observed MW: 43 kDa			
Applications	Tested Applications:		Positive Controls: WB : HL-60 cells, Raji cells, K-562 cells, Jurkat cells		
	WB, IHC, ELISA				
	Cited Applications: WB, IF, IP		IHC : mouse testis tissue, rat testis tissue		
	Species Specificity: human, mouse, rat				
	Cited Species: human, mouse				
		atively, antigen			
Background Information	human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GATA1 is first identified as a mer (WGATAR) binding motif [PMID:2 role in the normal development of confers transcriptional activity ar	atively, antigen d with citrate nber of the GATA trans 2937757]. GATA1, a zir of hematopoietic cell li nd a C-terminal domair	nc finger DNA-bin neages. The prote n that mediates b	nily, whose members bind the consens ding transcription factor, plays a critica ein contains an N-terminal region that inding to DNA and other factors [PMID: erythroid and megakaryocytic-specific	
	human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GATA1 is first identified as a mer (WGATAR) binding motif [PMID:2 role in the normal development of confers transcriptional activity ar 8524811]. GATA-1 is also implica genes [PMID:22937757].	atively, antigen d with citrate nber of the GATA trans 2937757]. GATA1, a zin of hematopoietic cell li nd a C-terminal domair ted in regulating the e	nc finger DNA-bin neages. The prote n that mediates b	ding transcription factor, plays a critica ein contains an N-terminal region that inding to DNA and other factors [PMID: rythroid and megakaryocytic-specific	
	human, mouse Note-IHC: suggested antiger TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GATA1 is first identified as a mer (WGATAR) binding motif [PMID:2 role in the normal development of confers transcriptional activity ar 8524811]. GATA-1 is also implica genes [PMID:22937757]. Author	atively, antigen d with citrate nber of the GATA trans 2937757]. GATA1, a zir of hematopoietic cell li nd a C-terminal domain ted in regulating the ex Pubmed ID Jo	nc finger DNA-bin neages. The prote n that mediates b xpression of the e	ding transcription factor, plays a critica ein contains an N-terminal region that inding to DNA and other factors [PMID:	
Background Information	human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GATA1 is first identified as a mer (WGATAR) binding motif [PMID:2 role in the normal development of confers transcriptional activity ar 8524811]. GATA-1 is also implica genes [PMID:22937757]. Author Wenqian Yu	atively, antigen d with citrate nber of the GATA trans 2937757]. GATA 1, a zir of hematopoietic cell li nd a C-terminal domair ted in regulating the er Pubmed ID Jo 31685800 Ce	nc finger DNA-bin neages. The proto n that mediates b xpression of the e urnal	ding transcription factor, plays a critical ein contains an N-terminal region that inding to DNA and other factors [PMID: rythroid and megakaryocytic-specific Application	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 60011-1-1g (GATA1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 60011-1-Ig (GATA1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).