### For Research Use Only

# STOML2 Monoclonal antibody

Catalog Number:60052-1-lg Featured Product

11 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 60052-1-lg BC002442 GeneID (NCBI): Concentration: 1600 ug/ml 30968 Source: **UNIPROT ID:** Q9UJZ1 Mouse

Isotype: Full Name: lgG2b stomatin (EPB72)-like 2

Immunogen Catalog Number: Calculated MW:

AG0363 356 aa, 39 kDa

> Observed MW: 39 kDa

**Purification Method:** Protein A purification

CloneNo.:

1A2E9

**Recommended Dilutions:** 

WB 1:5000-1:20000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000 IF/ICC 1:400-1:1600

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

**Cited Applications:** WB, IHC, IF, IP Species Specificity:

human, mouse, rat, pig

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval antigen retrieval with a trace buffer and 16.0 with citrate buffer pH 6.0

**Positive Controls:** 

WB: pig brain tissue, HepG2 cells, MCF-7 cells, RAW

264.7 cells

IP: mouse brain tissue.

IHC: human stomach cancer tissue, human

endometrial cancer tissue IF/ICC: A431 cells,

## **Background Information**

Human stomatin (band 7.2b) is a 31-kDa erythrocyte membrane protein of unknown function but implicated in the control of ion channel permeability, mechanoreception, and lipid domain organization. Stomatin (EPB72)-like 2 (STOML2, synonyms: SLP-2, HSPC108) is a 38.5-kDa protein that is overall approximately 20% similar to human stomatin. STOML2 is also present in mature human erythrocytes, but lacks a characteristic NH(2)-terminal  $hydrophobic\ domain\ found\ in\ other\ stomatin\ homologues.\ STOML2\ may\ link\ stomatin\ or\ other\ integral\ membrane$ proteins to the peripheral cytoskeleton and thereby play a role in regulating ion channel conductances or the organization of sphingolipid and cholesterol-rich lipid rafts.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Cheng-Ta Yang	29556045	Cell Death Dis	WB
Jingjing Zhang	30944651	Oncol Lett	IHC
Chongshu Jian	28630166	J Cell Sci	WB

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Pig, rat, and mouse brain tissues were subjected to SDS PAGE followed by western blot with 60052-1-Ig (STOML2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 60052-1-Ig (STOML2 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 60052-1-Ig (STOML2 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-STOML2 (IP:60052-1-1g, 4ug; Detection:60052-1-1g 1:1000) with mouse brain tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using STO ML2 antibody (60052-1-1g, Clone: 1A2E9) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).