For Research Use Only

## CXCR3B-specific Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

1B2D6

Catalog Number:60065-1-PBS

**Featured Product** 

**Basic Information** 

Catalog Number:

60065-1-PBS

Size: 1 mg/ml

Source: Mouse

IgG2b

GenBank Accession Number:

AF469635 GeneID (NCBI):

2833

UNIPROT ID: P49682 Full Name:

chemokine (C-X-C motif) receptor 3

Calculated MW: 46 kDa Observed MW:

43 kDa

**Applications** 

Tested Applications: WB, IHC, Indirect ELISA Species Specificity:

human

**Background Information** 

CXCR3 (C-X-C chemokine receptor type 3) is a G protein-coupled, seven-transmembrane domain chemokine receptor that is expressed on the surface of a number of cell types, including activated CD4+ and CD8+ T cells, NK and NK-T cells, plasmacytoid dendritic cells, and some B cells (PMID: 16847335). CXCR3 binds and is activated by the three IFN-gamma-inducible chemokines: CXCL9, CXCL10, and CXCL11. Binding of these ligands to CXCR3 has been implicated in regulating T-cell infiltration during inflammation and tissue injury (PMID: 17250586). Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms, CXCR3B, mediates the angiostatic activity of CXCR3 ligands and also acts as functional receptor for CXCL4 (PMID:12782716). This monoclonal antibody is specific to CXCR3B.

Storage

Storage

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


## **Selected Validation Data**



MOLT-4 cells were subjected to SDS PAGE followed by western blot with 60065-1-1g (CXCR3B-specific antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60065-1-Ig (CXCR3B-specific antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60065-1-Ig (CXCR3B-specific antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60065-1-PBS in a different storage buffer formulation.