For Research Use Only

RAC2 Monoclonal antibody, PBS Only

Catalog Number:60077-1-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

4C2H6

Basic Information

Catalog Number:

60077-1-PBS

Size: 1mg/ml

Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG1202

Full Name: ras-related C3 botulinum toxin

5880

P15153

BC001485

GeneID (NCBI):

UNIPROT ID:

substrate 2 (rho family, small GTP

GenBank Accession Number:

binding protein Rac2)

Calculated MW: 21 kDa

Observed MW: 21 kDa

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity: human, mouse, rat

Background Information

RAC2, also named as GX, belongs to the small GTPase superfamily and Rho family. RAC2 is a plasma membraneassociated small GTPase which cycles between an active GTP-bound and inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses, such as secretory processes, phagocytose of apoptotic cells and epithelial cell polarization. RAC2 augments the production of reactive oxygen species (ROS) by NADPH oxidase. Defects in RAC2 are the cause of neutrophil immunodeficiency syndrome.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60077-1-lg (RAC2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60077-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60077-1-lg (RAC2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60077-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60077-1-lg (RAC2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60077-1-PBS in a different storage buffer formulation.