For Research Use Only

GDI2 Monoclonal antibody

Catalog Number:60078-1-lg 1 Publications




Basic Information	Catalog Number: 60078-1-lg	GenBank Accession Number: BC005145 GeneID (NCBI): 2665 UNIPROT ID: P50395 Full Name: GDP dissociation inhibitor 2 og Number: Calculated MW: 47 kDa		Purification Method: Protein A purification	
	Size: 1000 µg/ml			CloneNo.: 4B8F3 Recommended Dilutions: WB 1:2000-1:10000 IHC 1:20-1:200 IF/ICC 1:400-1:1600	
	Source: Mouse				
	Isotype: IgG1				
	Immunogen Catalog Number: AG4613				
	Observed MW: 46 kDa		1W:		
Applications	IF/ICC, IHC, WB, ELISA WB : HeLa Cited Applications: cells, K-5 WB, IHC IHC : hum Species Specificity: tissue		Positive C	Controls:	
				.a cells, HepG2 cells, HEK-293 cells, Jurkat 562 cells, HSC-T6 cells, NIH/3T3 cells	
			an breast cancer tissue, human liver cancer		
	human, mouse, rat IF/ICC : HeLa cells, Cited Species:			eLa cells,	
	human Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	GDP dissociation inhibitors (GDIs) are proteins that regulate the GDP-GTP exchange reaction of members of the rab family, GDIs can bind and release GDP-bound Rab proteins from membranes. Two GDI proteins towards different Ral proteins have been identified. GDI 1 interacts with almost all of the Rab proteins, while GDI2 interacts with Rabll but not Rab3A. GDI2 distributes ubiquitously, displaying a membrane bound location in perinuclear regions of cells. GDI-2 was thought to be involved in cellular response to insulin. It electrophoreses as a 46kd protein in SDS-PAGE. (PMID: 7929030; PMID: 19570034). This antibody can bind both GDIs for the close sequences.				
Notable Publications	Author Pubmed ID Journal		Journal	Application	
	Zhiyue Ye	38911369	J Cancer	WB,IHC	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.1% sodium azide and Aliquoting is unnecessary for -20	I 50% glycerol p⊦			


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60078-1-Ig (GDI2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60078-1-1g (GDI2 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60078-1-1g (GDI2 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 60078-1-Ig (GDI2 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using GDI2 antibody (60078-1-lg, Clone: 4B8F3) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).