For Research Use Only

DGCR8 Monoclonal antibody

Catalog Number:60084-1-lg Featured Product

11 Publications



Basic Information

Applications

Catalog Number: 60084-1-lg Size: 1600 µg/ml

Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG4871

IP, WB, ELISA **Cited Applications:** WB, IP, RIP

Tested Applications:

Species Specificity: human, mouse, rat, pig Cited Species:

human, mouse, rat

GenBank Accession Number:

BC009323 GeneID (NCBI): 54487 **UNIPROT ID:** Q8WYQ5

DiGeorge syndrome critical region gene 8

Calculated MW: 773 aa, 86 kDa

Full Name:

Observed MW: 120 kDa

Positive Controls:

WB: HEK-293 cells, HepG2 cells, BxPC-3 cells, RAW 264.7 cells, A431 cells, Jurkat cells, K-562 cells,

Purification Method:

Protein A purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

WB 1:1000-1:6000

protein lysate

CloneNo.:

4D11A3

ROS1728 cells IP: K-562 cells,

Background Information

DGCR8 is a RNA-binding protein that assists the Rnase III enzyme Drosha in the processing of microRNAs (miRNAs), $which \ regulate \ the \ expression \ of \ a \ large \ number \ of \ protein-coding \ genes [PMID: 22580560]. \ DGCR8, \ which \ contains$ two double-stranded RNA (dsRNA)-binding domains, may be an essential component of the primary miRNAs processing complex, along with Drosha, promoting the processing of primary microRNA to precursor microRNA. It is ubiquitous expressed in human and mouse tissues, and is deleted in DiGeorge syndrome[22323604]. The calculated molecular weight of DGCR8 is 82-86 kDa, but the post-modified DGCR8 is about 120 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Haidong Xu	34528746	Aging Cell	WB
Hailong Zhang	34518544	Nat Commun	WB
Fangfang Zhang	31813799	Mol Ther	WB,RIP


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60084-1-Ig (DGCR8 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-DGCR8 (IP:60084-1-lg, 5ug; Detection:60084-1-lg 1:500) with K-562 cells lysate 3440ug.