

IgA Monoclonal antibody

Catalog Number: 60099-1-Ig 2 Publications

Basic Information

Catalog Number: 60099-1-Ig	GenBank Accession Number: BC016369	Purification Method: Caprylic acid/ammonium sulfate precipitation
Size: 1900 μ g/ml	GeneID (NCBI): 3493	CloneNo.: 3A5A5
Source: Mouse	Full Name: immunoglobulin heavy constant alpha 1	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:160000-1:640000 IF 1:50-1:500
Isotype: IgG1	Calculated MW: 496 aa, 53 kDa	
Immunogen Catalog Number: AG7417	Observed MW: 55-60 kDa	

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA	Positive Controls:
Cited Applications: IF	WB : human saliva tissue, human saliva, human plasma
Species Specificity: human	IHC : human tonsillitis tissue,
Cited Species: rat	IF : human tonsillitis tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Ig alpha is the major immunoglobulin class in body secretions. It may serve both to defend against local infection and to prevent access of foreign antigens to the general immunologic system. This antibody recognizes the Chain C region of human Ig alpha-1 (IGHA1).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhipeng Li	30670042	J Nanobiotechnology	
Meiying Chang	37972847	Eur J Pharmacol	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human saliva were subjected to SDS PAGE followed by western blot with 60099-1-Ig (Human IgA Antibody) at dilution of 1:16000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60099-1-Ig (Human IgA antibody) at dilution of 1:320000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 60099-1-Ig (Human IgA antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).