For Research Use Only

NR2F6 Monoclonal antibody, PBS Only

Concentration: 1mg/ml

Catalog Number:60117-2-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

2H2B8

Basic Information

Catalog Number: 60117-2-PBS

BC002669

GenBank Accession Number:

GeneID (NCBI): 2063

Source: UNIPROT ID: Mouse P10588
Isotype: Full Name:

IgG1 nuclear receptor subfamily 2, group F,

Immunogen Catalog Number:

AG7614 Calculated MW:

43 kDa

Observed MW: 43 kDa

Applications

Tested Applications: WB, IHC, Indirect ELISA Species Specificity: human, rat

Background Information

NR2F6 belongs to NR2F subfamily members, which have their roles in the regulation of neurogenesis, organogenesis, and cellular differentiation during embryonic development (PMID:15875024). NR2F6 acts as transcription factor by binding to a TGACCT direct-repeat motif and has been established in controlling facets of central nervous system (CNS) (PMID: 12690040). It is a nuclear attenuator that directly interferes with DNA binding of NF-AT and, subsequently, transcriptional activity of the NF-AT-dependent IL-17 expression. Otherwise, NR2F6 negatively interfered with transcriptional cytokine responses and acted as a barrier against autoimmunity (PMID:18701084).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C $\,$

Storage Buffer: PBS Only


Selected Validation Data



WB result of NR2F6 antibody (60117-2-Ig, 1:1000) with si-Control and si-NR2F6 transfected HeLa cells. This data was developed using the same antibody clone with 60117-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60117-2-Ig (NR2F6 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60117-2-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 60117-2-lg (NR2F6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60117-2-PBS in a different storage buffer formulation.