For Research Use Only

CD206 Monoclonal antibody

Catalog Number:60143-1-lg Featured Product 205 Publications



Basic Information	Catalog Number: 60143-1-lg	GenBank Ac NM_002438	GenBank Accession Number: NM_002438		Purification Method: Protein A purification	
	Concentration:GeneID (NCBI):2000 ug/ml4360		BI):	CloneNo.:		
			2A6A10			
	Source:	UNIPROT ID:		Recommended Dilutions:		
	Mouse	Full Name:	Full Name: mannose receptor, C type 1 Calculated MW: 166 kDa		WB: 1:5000-1:50000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC: 1:10000-1:40000 IF-P: 1:200-1:800	
	IgG2a	mannose re				
	-	Calculated 166 kDa				
	Observed MW: 170 kDa		IW:			
Applications	Tested Applications:		Positive Controls:			
	WB, ITC, IF-P, IP, ELISA WB : hum		an placenta tissue, human liver tissue			
	WB, IHC, IF		IP : human placenta tissue,			
	Species Specificity: human		IHC : human lung cancer tissue, human liver tissue, human placenta tissue			
	Cited Species: IF-P : human lung cancer tissue, human, pig					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	CD206, also named as MMR, CLEC13D and MRC1, is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. CD206 has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. CD206 is a 170 kDa transmembrane protein which contains 5 domains: an amino-terminal cysteine-rich region, a fibronectin type II repeat, a series of eight tandem lectin-like carbohydrate recognition domains (responsible for the recognition of mannose and fucose), a transmembrane domain, and an intracellular carboxy-terminal tail. It is expressed on most tissue macrophages, in vitro derived dendritic cells, lymphatic and sinusoidal endothelia. This antibody recognizes the intracellular carboxy-terminal part of CD206 and MRC1L1.					
Notable Publications	Author	Pubmed ID	Journal		Application	
	Xinmei Huang	34478541	J Clin Endocrinol I	Metab	IHC	
	Liping Xu	36179453	Tissue Cell		IHC	
	C. Zhao	34647005	Mater Today Bio		IF	
Storage	Storage: Store at -20°C. Stable for on Storage Buffer: PBS with 0.02% sodium azi Aliquoting is unnecessary f	e year after shipment. de and 50% glycerol, p for -20 [°] C storage	H7.3			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





human placenta tissue was subjected to SDS PAGE followed by western blot with 60143-1-lg (CD206 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 60143-1-lg (CD206 antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 60143-1-1g (CD206 antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using CD206 antibody (60143-1-Ig, Clone: 2A6A10) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using CD206 antibody (60143-1-Ig, Clone: 2A6A10) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP result of anti-CD206 (IP:60143-1-Ig, 5ug; Detection:60143-1-Ig 1:300) with human placenta tissue lysate 1520ug.