For Research Use Only

CD109 Monoclonal antibody

Catalog Number:60179-1-lg 3 Publications



Basic Information

Catalog Number: 60179-1-lg Size: 1300 µg/ml Source:

Mouse Isotype: IgG1 GenBank Accession Number:

GenBank Accession Nur BC152996 GeneID (NCBI): 135228 UNIPROT ID: Q6YHK3 Full Name: CD109 molecule Calculated MW: 162 kDa Observed MW:

162 kDa

Purification Method:

Caprylic acid/ammonium sulfate precipitation

CloneNo.: 2H8G7

Recommended Dilutions: WB 1:1000-1:8000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human
Cited Species:
human

Positive Controls:

WB: Jurkat cells,

Background Information

CD109, also named as CPAMD7, is a member of the alpha2-macroglobulin/complement superfamily. It is a glycosylphosphatidylinositol (GPI)-linked cell surface glycoprotein of approximately 170kd which is found on the cell surface of platelets, activated T-cells, and endothelial cells. CD109 binds to and negatively regulates signaling of transforming growth factor beta (TGF-beta). CD109 has been identified as part of the TGF-beta receptor system in human keratinocytes. Up regulation of CD109 expression has been observed in several different types of tumor.CD109 is a glycosyl-phosphatidylinositol-anchored glycoprotein that is a member of the a2-macroglobulin/C3, C4, C5 family of thioester-containing proteins. There're 2 forms of 150 (p150) and 120 kDa (p120) exist due to proteolytic degradation from a 180 kDa form. The antibody can recognize all these 4 isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Tobias Fretwurst	31877027	Clin Oral Implants Res	WB
Junyu Fan	31403228	Drug Dev Res	
Yunyan Liu	37378424	Mol Carcinog	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 60179-1-1g (CD109 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.