

For Research Use Only

CD138/Syndecan-1 Monoclonal antibody



Catalog Number: 60185-2-Ig **1 Publications**

Basic Information

Catalog Number: 60185-2-Ig	GenBank Accession Number: BC008765	Purification Method: Protein A purification
Size: 600 µg/ml	GeneID (NCBI): 6382	CloneNo.: 4H5H5
Source: Mouse	UNIPROT ID: P18827	Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200
Isotype: IgG2a	Full Name: syndecan 1	
Immunogen Catalog Number: AG0823	Calculated MW: 32.5 kDa	
	Observed MW: 140 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : MCF7 cells, IHC : human tonsil tissue,
Cited Applications: WB, ColP	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Syndecan-1 (SDC1), also named CD138, is an integral membrane protein. It participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. It is a heparan sulfate proteoglycan expressed on the surface of, and actively shed by, myeloma cells. Altered syndecan-1 expression has been detected in several different tumor types. SDC1 was regarded as a useful marker for labeling normal and neoplastic plasma cells and plasmacytoid lymphomas.

Notable Publications

Author	Pubmed ID	Journal	Application
Charlie Colin-Pierre	34957110	Front Cell Dev Biol	WB, ColP

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



MCF7 cells were subjected to SDS PAGE followed by western blot with 60185-2-Ig (CD138/Syndecan-1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsil using 60185-2-Ig (CD138/Syndecan-1 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human tonsil using 60185-2-Ig (CD138/Syndecan-1 antibody) at dilution of 1:50 (under 40x lens).