## For Research Use Only

## WTAP Monoclonal antibody

Catalog Number:60188-1-lg

Featured Product

109 Publications



**Basic Information** 

Catalog Number: 60188-1-lg Source: Mouse

Isotype: IgG1

Immunogen Catalog Number:

AG0259

GenBank Accession Number:

BC000383 GeneID (NCBI): 9589 UNIPROT ID: Q15007

Full Name: Wilms tumor 1 associated protein

Calculated MW: 45 kDa Observed MW: 50 kDa Purification Method:

Protein G purification

CloneNo.: 4A10G9

Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:500-1:2000 IF/ICC: 1:50-1:500

**Applications** 

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, RIP Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat, drosophila

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Positive Controls:** 

WB: MCF-7 cells, HeLa cells, HEK-293 cells, HepG2 cells, LNCaP cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells

IP: HeLa cells,

IHC: human colon tissue,
IF/ICC: HeLa cells,

**Background Information** 

WTAP, also named as KIAAO105 and hFL(2)D, is previously identified as a protein associated with Wilms' tumor-1 (WT-1) protein that is essential for the development of the genitourinary system. It is a Pre-mRNA-splicing regulator protein. It is a regulatory subunit of the WMM N6-methyltransferase complex which plays a role in the efficiency of mRNA splicing and processing and mRNA stability. It is required for accumulation of METTL3 and METTL14 to nuclear speckle. MW of WTAP is 45-50 kDa due to phosphorylation.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Fanpeng Zhao	34593014	Mol Neurodegener	WB
Jing Wang	32998774	J Ovarian Res	WB
Qian Luo	36175708	Med Oncol	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60188-1-Ig (WTAP antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-WTAP (IP:60188-1-Ig, 4ug; Detection:10200-1-AP 1:8000) with HeLa cells lysate 2000 ug.



Immunofluorescent analysis of (4% PFA ) fixed HeLa cells using 60188-1-1g(WTAP antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 60188-1-lg (WTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 60188-1-lg (WTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.