For Research Use Only

PPARD Monoclonal antibody

Catalog Number:60193-1-lg Featured Product

10 Publications



Basic Information

Applications

Catalog Number: 60193-1-lg Concentration: 740 ug/ml Source:

Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG16770

Tested Applications:

WB, ELISA Cited Applications:

WB, ChIP

Species Specificity: human, pig **Cited Species:**

human, mouse, rat, zebrafish

GenBank Accession Number:

BC002715 GeneID (NCBI):

UNIPROT ID: Q03181

5467

Full Name: peroxisome proliferator-activated

receptor delta

Calculated MW: 50 kDa Observed MW:

54 kDa

WB: HeLa cells, A549 cells, pig adipose tissue, Jurkat

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:1000-1:6000

CloneNo.:

1B10E1

cells, K-562 cells

Positive Controls:

Background Information

Peroxisome proliferator-activator receptor delta(PPARD), a ligand-activated transcription factor, belongs to the $PPARs\ family, which\ get its\ name\ for\ their\ chemicals\ that\ induce\ proliferation\ of\ peroxisomes,\ organelles\ that$ contributes to the oxidation of fatty acids. PPARD is a receptor for peroxisome proliferator and once activated by a ligand, it binds to and activates the promoter elements of target genes, such as acylCoA oxidase gene. But it act as a repressor for the NPC1L1 gene.

Notable Publications

Author	Pubmed ID	Journal	Application
Alyssa Charrier	27624101	Am J Physiol Endocrinol Metab	WB
Yunzhi Wang	29674717	Nat Commun	WB
Xiaolong Qu	39862078	Front Biosci (Landmark Ed)	WB


Storage


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 60193-1-1g (PPARD antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



pig adipose tissue were subjected to SDS PAGE followed by western blot with 60193-1-1g (PPARD antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.