For Research Use Only

Fas/CD95 Monoclonal antibody

Catalog Number:60196-1-lg Featured Product 15 Publications


oroteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 60196-1-lg	GenBank Accession Numb BC012479	er: Purification Method: Protein G purification	
	Concentration:	GenelD (NCBI):	CloneNo.:	
	1000 ug/ml	355	1G7F7	
	Source:	UNIPROT ID:	Recommended Dilutions:	
	Mouse P25445 Isotype: Full Name:		WB: 1:1000-1:8000	
			IHC: 1:50-1:500	
	lgG1	Fas (TNF receptor superfamily, member 6) Calculated MW:		
	Immunogen Catalog Number: AG3718			
		35-38 kDa		
		Observed MW: 38-45 kDa		
Applications	Tested Applications:	Po	Positive Controls:	
			3 : Jurkat cells, HeLa cells, K-562 cells, U-937 cells	
	Cited Applications: WB, IHC, IF		IHC : human tonsillitis tissue, human colon cancer tissue	
	Species Specificity: human			
	Cited Species: human, zebrafish			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	retrieval may be performed			
Background Information	retrieval may be performed buffer pH 6.0 FAS, also named as CD95, APO-1, receptor belonging to the TNF rece antibody or Fas ligand. Stimulatio the formation of the death-inducin induction of peripheral tolerance,	APT1, FAS1 and TNFRSF6, is a eptor superfamily, can mediate on of Fas results in the aggrega ng signaling complex (DISC). F in the antigen-stimulated suic his anti-Fas monoclonal antibo	tion of its intracellular death domains, leading to AS-mediated apoptosis may have a role in the ide of mature T-cells, or both. The secreted isofor	
Background Information	retrieval may be performed buffer pH 6.0 FAS, also named as CD95, APO-1, receptor belonging to the TNF rece antibody or Fas ligand. Stimulatio the formation of the death-inducin induction of peripheral tolerance, 2 to 6 block apoptosis (in vitro). The through Fas by imitating the Fas-	APT1, FAS1 and TNFRSF6, is a eptor superfamily, can mediate on of Fas results in the aggrega ng signaling complex (DISC). F in the antigen-stimulated suic his anti-Fas monoclonal antibo ligand.	e apoptosis by ligation with an agonistic anti-Fas tion of its intracellular death domains, leading to AS-mediated apoptosis may have a role in the ide of mature T-cells, or both. The secreted isofor dy can be used to induce apoptosis in cell culture	
	retrieval may be performed buffer pH 6.0 FAS, also named as CD95, APO-1, receptor belonging to the TNF rece antibody or Fas ligand. Stimulatic the formation of the death-inducin induction of peripheral tolerance, 2 to 6 block apoptosis (in vitro). The through Fas by imitating the Fas-In- Author	APT1, FAS1 and TNFRSF6, is a eptor superfamily, can mediate on of Fas results in the aggrega ng signaling complex (DISC). F in the antigen-stimulated suic his anti-Fas monoclonal antibo ligand.	e apoptosis by ligation with an agonistic anti-Fas tion of its intracellular death domains, leading to AS-mediated apoptosis may have a role in the ide of mature T-cells, or both. The secreted isofor dy can be used to induce apoptosis in cell culture Application	
Background Information	retrieval may be performed buffer pH 6.0 FAS, also named as CD95, APO-1, receptor belonging to the TNF rec antibody or Fas ligand. Stimulatio the formation of the death-induci induction of peripheral tolerance, 2 to 6 block apoptosis (in vitro). TI through Fas by imitating the Fas- Author Liyuan Ran	APT1, FAS1 and TNFRSF6, is a eptor superfamily, can mediate on of Fas results in the aggrega ng signaling complex (DISC). F in the antigen-stimulated suic his anti-Fas monoclonal antibo ligand.	e apoptosis by ligation with an agonistic anti-Fas tion of its intracellular death domains, leading to AS-mediated apoptosis may have a role in the ide of mature T-cells, or both. The secreted isofor dy can be used to induce apoptosis in cell culture Application WB	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Jurkat cells were subjected to SDS PAGE followed by western blot with 60196-1-1g (FAS/CD95 antibody at dilution of 1:4000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60196-1-1g (FAS/CD95 antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60196-1-1g (FAS/CD95 antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).