## For Research Use Only

## p504S,AMACR Monoclonal antibody

Catalog Number: 60240-1-lg



**Basic Information** 

Catalog Number: GenBank Accession Number:

60240-1-lg BC009471

Size: GeneID (NCBI):
930 ug/ml 23600

Source: UNIPROT ID:
Mouse Q9UHK6

Isotype: Full Name:

IgG1 alpha-methylacyl-CoA racemase

Immunogen Catalog Number:Calculated MW:AG8720382 aa, 42 kDaObserved MW:

42 kDa

**Applications** 

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

human, mouse, rat

**Positive Controls:** 

WB: HepG2 cells, PC-3 cells, HEK-293 cells, HeLa cells, LNCaP cells, HSC-T6 cells, 4T1 cells, NIH/3T3 cells

**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:1000-1:6000 IF/ICC 1:200-1:800

CloneNo.:

2H9B5

IF/ICC: A431 cells,

## **Background Information**

AMACR(Alpha-methyl acyl-CoA racemase) belongs to the CaiB/BaiF CoA-transferase family. It is a mitochondrial and peroxisomal enzyme that catalyzes the conversion of 2R stereoisomers of phytanic and pristanic acid to their S counterparts. AMACR has previously been shown to be a highly sensitive marker for colorectal and clinically localized prostate cancer (PCa). However, AMACR expression is down-regulated at the transcript and protein level in hormone-refractory metastatic PCa, suggesting a hormone-dependent expression of AMACR(PMID:12213712). It has 3 isoforms produced by alternative splicing. Defects in AMACR are the cause of alpha-methyl acyl-CoA racemase deficiency (AMACRD) and congenital bile acid synthesis defect type 4 (CBAS4).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 60240-1-1g (p504S,AMACR antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 60240-1-1g (p504S,AMACR antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using p504S,AMACR antibody (60240-1-lg, Clone: 2H9B5) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).