For Research Use Only

CD14 Monoclonal antibody

Catalog Number:60253-1-lg 18 Publications



Basic Information

Catalog Number: 60253-1-lg Concentration:

1000 ug/ml
Source:
Mouse
Isotype:
IgG1

Immunogen Catalog Number:

AG10693

GenBank Accession Number: BC010507

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000170458 UNIPROT ID: P08571 Full Name:

CD14 molecule

Calculated MW: 375 aa, 40 kDa Observed MW: 45-55 kDa Purification Method:

Protein G purification

CloneNo.: 2C1D9

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:1000-1:4000 IF-P 1:50-1:500 IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse Cited Species:

human, mouse, rabbit

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: THP-1 cells, human milk, human placenta tissue,

RAW 264.7 cells

IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,

IF/ICC: RAW 264.7 cells, human tonsillitis tissue

Background Information

CD14 is a 50-55 kDa glycosylphosphatidylinositol-anchored glycoprotein preferentially expressed on monocytes and macrophages, and at lower levels on granulocytes (PMID: 3385210; 2462937; 7685797). CD14 can also exist as a soluble protein. CD14 acts as a co-receptor for bacterial liposaccharides (LPS) (PMID: 1698311). It plays a major role in the inflammatory response of monocytes to LPS.

Notable Publications

Author	Pubmed ID	Journal	Application
Zeqing Zhai	34480835	Arthritis Rheumatol	IF
Haochen Hui	36208807	Int J Biol Macromol	WB
Jia Xiong	35320754	Mol Cell	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




THP-1 cells were subjected to SDS PAGE followed by western blot with 60253-1-Ig (CD14 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours




Untreated and PNGase F-treated lysates of human milk were subjected to SDS PAGE followed by western blot with 60253-1-Ig (CD14 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-Ig (CD14 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-1g (CD14 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD14 antibody (60253-1-lg, Clone: 2C1D9) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of RAW 264.7 cells using 60253-1-Ig (CD14 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).