For Research Use Only

CD14 Monoclonal antibody, PBS Only (Capture)

Catalog Number: 60253-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein G purification

60253-1-PBS Size:

GeneID (NCBI):

CloneNo.:

1000 $\,\mu$ g/ml

BC010507

2C1D9

Source: Mouse Isotype: lgG1

ENSG00000170458 UNIPROT ID: P08571

ENSEMBL Gene ID:

Immunogen Catalog Number: AG10693

Full Name: CD14 molecule

Calculated MW: 375 aa, 40 kDa Observed MW:

45-55 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, Cytometric bead array, Indirect

Species Specificity:

human, mouse

Background Information

 $CD14 is \ a \ 50-55 \ kDa \ glycosylphosphatidy linositol-anchored \ glycoprotein \ preferentially \ expressed \ on \ monocytes$ and macrophages, and at lower levels on granulocytes (PMID: 3385210; 2462937; 7685797). CD14 can also exist as a soluble protein. CD14 acts as a co-receptor for bacterial liposaccharides (LPS) (PMID: 1698311). It plays a major role in the inflammatory response of monocytes to LPS.


Storage

Storage:


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only

Selected Validation Data




THP-1 cells were subjected to SDS PAGE followed by western blot with 60253-1-lg (CD14 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.




Untreated and PNGase F-treated lysates of human milk were subjected to SDS PAGE followed by western blot with 60253-1-Ig (CD14 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-1g (CD14 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-lg (CD14 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD14 antibody (60253-1-lg, Clone: 2C1D9) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of RAW 264.7 cells using 60253-1-19 (CD14 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Affini Pure Goat Anti-Mouse IgG (H+L). This data was developed using the same antibody clone with 60253-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP50282-2, CD14 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 60253-1-PBS. Detection antibody: 60253-2-PBS. Standard:Ag10693. Range: 1.563-200 ng/mL