

For Research Use Only

SMN-Exon7 Monoclonal antibody

Catalog Number: 60255-1-Ig **1 Publications**



Basic Information

Catalog Number: 60255-1-Ig	GenBank Accession Number: BC062723	Purification Method: Protein A purification
Concentration: 1000 ug/ml	GeneID (NCBI): 6606	CloneNo.: 3A8G11
Source: Mouse	UNIPROT ID: Q16637	Recommended Dilutions: WB 1:500-1:2000 IHC 1:1000-1:4000 IF/ICC 1:50-1:500
Isotype: IgG1	Full Name: survival of motor neuron 1, telomeric	
Immunogen Catalog Number: AG16615	Calculated MW: 294 aa, 32 kDa Observed MW: 40 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : HEK-293 cells, HeLa cells, HepG2 cells IHC : mouse heart tissue, mouse brain tissue, mouse pancreas tissue, rat brain tissue, rat heart tissue, rat pancreas tissue IF/ICC : HepG2 cells,
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Spinal muscular atrophy (SMA) is an autosomal recessive neurodegenerative disease characterized by loss of anterior horn cells in the spinal cord and concomitant symmetrical muscle weakness and atrophy (PMID: 16364894). SMA is caused by deletion or mutations of the survival motor neuron (SMN1) gene. SMA patients lack a functional SMN1 gene, but they possess an intact SMN2 gene, which though nearly identical to SMN1, is only partially functional (PMID: 17355180). A large majority of SMN2 transcripts lack exon 7, resulting in production of a truncated, less stable SMN protein (PMID: 10369862). The level of SMN protein correlates with phenotypic severity of SMA. This antibody, 60255-1-Ig, raised against the C-terminal region (275-294aa) encoded by the exon 7.

Notable Publications

Author	Pubmed ID	Journal	Application
Mandana Arbab	36996170	Science	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat heart tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat pancreas tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).