For Research Use Only

SMN-Exon7 Monoclonal antibody, PBS Only




Catalog Number:60255-1-PBS


| Basic Information | Catalog Number: 60255-1-PBS | GenBank Accession Number: BC062723 | Purification Method: Protein A purification |
|------------------------|--|---------------------------------------|--|
| | Size: 1mg/ml | GenelD (NCBI): 6606 | CloneNo.: 3A8G11 |
| | Source: Mouse | UNIPROT ID: Q16637 | |
| | Isotype:Full Name:IgG1survival of motor neuron 1, telomeric | | neric |
| | Immunogen Catalog Number: AG16615 | Calculated MW: 294 aa, 32 kDa | |
| | | Observed MW: 40 kDa | |
| Applications | Tested Applications: WB, IHC, IF/ICC, Indirect ELISA | | |
| | Species Specificity: human, mouse, rat | | |
| Background Information | Spinal muscular atrophy (SMA) is an autosomal recessive neurodegenerative disease characterized by loss of anterior horn cells in the spinal cord and concomitant symmetrical muscle weakness and atrophy (PMID: 16364894). SMA is caused by deletion or mutations of the survival motor neuron (SMN1) gene. SMA patients lack a functional SMN1 gene, but they possess an intact SMN2 gene, which though nearly identical to SMN1, is only partially functional (PMID: 17355180). A large majority of SMN2 transcripts lack exon 7, resulting in production of a truncated, less stable SMN protein (PMID: 10369862). The level of SMN protein correlates with phenotypic severity of SMA. This antibody, 60255-1-lg, raised against the C-terminal region (275-294aa) encoded by the exon 7. | | |
| Storage | Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only | | |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data









Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse pancreas tissue slide using 60255-1-1g (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 60255-1-lg (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat heart tissue slide using 60255-1-lg (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat pancreas tissue slide using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of HepG2 cells using 60255-1-Ig (SMN-Exon7 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L). This data was developed using the same antibody clone with 60255-1-PBS in a different storage buffer formulation.