

For Research Use Only

CD11c/Integrin Alpha X Monoclonal antibody, PBS Only



Catalog Number: 60258-1-PBS

Basic Information

Catalog Number: 60258-1-PBS	GenBank Accession Number: BC038237	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 3687	CloneNo.: 2F1C10
Source: Mouse	ENSEMBL Gene ID: ENSG00000140678	
Isotype: IgG2a	UNIPROT ID: P20702	
Immunogen Catalog Number: AG11350	Full Name: integrin, alpha X (complement component 3 receptor 4 subunit)	
	Calculated MW: 1169 aa, 129 kDa	
	Observed MW: 145 kDa	

Applications

Tested Applications:
WB, IHC, IF, ELISA
Species Specificity:
human

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits. CD11c/Integrin alpha X is a 145-150 kDa type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, NK cells and dendritic cells. Integrin alpha X/beta 2 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C
Storage Buffer:
PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Untreated and PNGase F-treated lysates of HL-60 cells were subjected to SDS PAGE followed by western blot with 60258-1-Ig (CD11c/Integrin alpha X antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808). This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.




human spleen tissue were subjected to SDS PAGE followed by western blot with 60258-1-Ig (CD11c/Integrin alpha X Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60258-1-Ig (CD11c/Integrin Alpha X antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60258-1-Ig (CD11c/Integrin Alpha X antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD11c/Integrin Alpha X antibody (60258-1-Ig, Clone: 2F1C10) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD11c/Integrin Alpha X antibody (60258-1-Ig, Clone: 2F1C10) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60258-1-PBS in a different storage buffer formulation.