

For Research Use Only

P53 Monoclonal antibody

Catalog Number: 60283-2-Ig

Featured Product

276 Publications



Basic Information

Catalog Number:

60283-2-Ig

Concentration:

1000 ug/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG0698

GenBank Accession Number:

BC003596

GeneID (NCBI):

7157

UNIPROT ID:

P04637

Full Name:

tumor protein p53

Calculated MW:

44 kDa

Observed MW:

53 kDa

Purification Method:

Protein A purification

CloneNo.:

6C4B6

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:1600-1:6400

IF/ICC 1:400-1:1600

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP

Species Specificity:

human, rat

Cited Species:

human, rat, pig, rabbit, canine, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A431 cells, COLO 320 cells, HSC-T6 cells, ROS1728 cells, rat thymus tissue, HaCaT cells, HT-29 cells, HEK-293 cells

IHC : human colon cancer tissue, human breast cancer tissue, human gliomas tissue, human lung cancer tissue, human ovary cancer tissue, human ovary tumor tissue, human stomach cancer tissue, human tonsillitis tissue

IF/ICC : A431 cells,

Background Information

TP53, also known as P53 and NY-CO-13, belongs to the p53 family and has 9 isoforms. In SDS-Page, the observed molecular weight is about 53 kDa. TP53 acts as a tumor suppressor in many tumor types, including growth arrest or apoptosis depending on the physiological circumstances and cell types. It is involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. TP53 Localizes in the nucleus in most cells but found in the cytoplasm in some cells. (PMID: 26166714; PMID: 25225161)

Notable Publications

Author	Pubmed ID	Journal	Application
Jie Liu	36238648	Oxid Med Cell Longev	WB
Xiaofang Zhu	36174712	Life Sci	WB
Yingjie Qing	34603598	Oxid Med Cell Longev	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60283-2-Ig (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 60283-2-Ig (P53 antibody) at dilution of 1:3200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




HSC-T6 cells were subjected to SDS PAGE followed by western blot with 60283-2-Ig (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




ROS1728 cells were subjected to SDS PAGE followed by western blot with 60283-2-Ig (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



rat thymus tissue were subjected to SDS PAGE followed by western blot with 60283-2-Ig (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 60283-2-Ig (P53 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using P53 antibody (60283-2-Ig, Clone: 6C4B6) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red).