For Research Use Only

ATP5C1 Monoclonal antibody, PBS Only



Catalog Number: 60284-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number: BC000931

Purification Method:

60284-1-PBS

GeneID (NCBI):

Protein A purification

Size: 1mg/ml

eneID (NCBI)

CloneNo.:

1mg/ml Source:

UNIPROT ID: P36542

1B5A2

Mouse Isotype: IgG2a

AG1343

Full Name: ATP synthase, H+ transporting,

Immunogen Catalog Number:

mitochondrial F1 complex, gamma

polypeptide 1

Calculated MW:

33 kDa

Observed MW:

33 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human

Background Information

ATP5C1(ATP synthase subunit gamma, mitochondrial) is also named as ATP5C, ATP5CL1 and belongs to the ATPase gamma chain family. Some scientists reported the complete sequence of the gene for the human ATP synthase gamma subunit and described tissue-specific isoforms of the subunit generated by alternative splicing of exon 9. The liver (L) isoform differed from the heart (H) isoform by the addition of a single amino acid (asp273) at the C terminus. (PMID:8227057).

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


Selected Validation Data




human heart tissue were subjected to SDS PAGE followed by western blot with 60284-1-lg (ATP5C1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60284-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60284-1-Ig (ATP5C1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60284-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60284-1-Ig (ATP5C1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60284-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 60284-1-lg (ATP5C1 antibody), at dilution of 1:100 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60284-1-PBS in a different storage buffer formulation



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human ATPSC1 (60284-1-ig, Clone:1B5A2) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 60284-1-PBS in a different storage buffer formulation.