For Research Use Only

PRDX4 Monoclonal antibody

Catalog Number:60286-1-lg Featured Product

3 Publications




Basic Information	Catalog Number: 60286-1-lg	GenBank Accession N BC007107	ank Accession Number: 7107		Purification Method: Protein A purification	
	Size:	GeneID (NCBI):		CloneNo.: 5B6F1		
	1200 µg/ml	10549				
	Source: Mouse	UNIPROT ID: Q13162		Recommended Dilutions: WB 1:5000-1:50000		
	lsotype: lgG2b	Full Name: peroxiredoxin 4		IHC 1:20-1:200 IF/ICC 1:400-1:1600		
	Immunogen Catalog Number: AG13401	Calculated MW: 31 kDa				
		Observed MW: 30 kDa				
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA					
	WD: LINCAP CEILS,		.ls, NIH/3T3 cells, HeLa cells, HepG2 lls, PC-12 cells, HEK-293 cells, Jurkat :ells			
				: human colon tissue, human pancreas cancer ie, human prostate cancer tissue		
	Cited Species: human, mouse		IF/ICC : HepC	IF/ICC : HepG2 cells,		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
	PRDX4(Peroxiredoxin-4) is also named as AOE37-2, Prx-IV and belongs to the AhpC/TSA family. PRDX4 is associate with acrosome formation during rat spermatogenesis and has a protective role in the male reproductive tract, because the phenotype of mice lacking this isoform includes testicular atrophy and increased sperm DNA damage(PMID:20864641). It is initially synthesized as a membrane-binding 31-kDa protein and processed into a 27 kDa secretory form and is discarded with the residual bodies(PMID:19208552). PRDX4 is a pentamer of dimers(PMID:23025503).					
Background Information	with acrosome formation during because the phenotype of mice la damage(PMID:20864641). It is in kDa secretory form and is discard	at spermatogenesis and h acking this isoform include tially synthesized as a me	es testicular atro embrane-bindin	pphy and increased g 31-kDa protein a	productive tract, sperm DNA nd processed into a 2	
	with acrosome formation during because the phenotype of mice la damage(PMID:20864641). It is in kDa secretory form and is discard	at spermatogenesis and h acking this isoform include tially synthesized as a me	es testicular atro embrane-bindin es(PMID:192085)	pphy and increased g 31-kDa protein a	productive tract, sperm DNA nd processed into a 2	
	with acrosome formation during because the phenotype of mice la damage(PMID:20864641). It is in kDa secretory form and is discard dimers(PMID:23025503).	rat spermatogenesis and h acking this isoform include tially synthesized as a me ed with the residual bodie Pubmed ID Journ	es testicular atro embrane-bindin es(PMID:192085)	ophy and increased g 31-kDa protein a 52). PRDX4 is a per	oroductive tract, sperm DNA nd processed into a 2 tamer of	
	with acrosome formation during because the phenotype of mice la damage(PMID:20864641). It is in kDa secretory form and is discard dimers(PMID:23025503).	rat spermatogenesis and h acking this isoform include tially synthesized as a me ed with the residual bodie Pubmed ID Journ 36334439 Bioch	es testicular atro embrane-bindin es(PMID:192085)	ophy and increased g 31-kDa protein a 52). PRDX4 is a per	oroductive tract, sperm DNA nd processed into a 2 tamer of Application	
Background Information	with acrosome formation during because the phenotype of mice la damage(PMID:20864641). It is in kDa secretory form and is discard dimers(PMID:23025503). Author Wangjuan Dai	rat spermatogenesis and h acking this isoform include tially synthesized as a me ed with the residual bodie Pubmed ID Journ 36334439 Bioch 36357400 Nat C	es testicular atro embrane-bindin es(PMID:192085) nal em Biophys Res	ophy and increased g 31-kDa protein a 52). PRDX4 is a per	oroductive tract, sperm DNA nd processed into a z tamer of Application IF	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of PRDX4 antibody (60286-1-lg, 1:5000) with si-Control and si-PRDX4 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 60286-1-lg (PRDX4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon slide using 60286-1-lg (PRDX4 Antibody) at dilution of 1:50(unde10x lens).



Immunohistochemical analysis of paraffinembedded human colon slide using 60286-1-1g (PRDX4 Antibody) at dilution of 1:50(under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PRDX4 antibody (60286-1-Ig, Clone: 5B6F1) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).