For Research Use Only

PLAP Monoclonal antibody, PBS Only

Catalog Number:60294-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number: BC009647

Purification Method: Protein G purification

60294-1-PBS

GeneID (NCBI):

Size: 1mg/ml

250

CloneNo.: 2B7C9

Source: Mouse

UNIPROT ID: P05187 Full Name:

Isotype: lgG1

alkaline phosphatase, placental

Immunogen Catalog Number:

(Regan isozyme)

AG13361

Calculated MW:

58 kDa

Observed MW:

66 kDa

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human

Background Information

 $Placental\ alkaline\ phosphatase\ (PLAP)\ is\ a\ glycosyl\ phosphatidylinositol\ (GPI)-anchored\ sialoglycoprotein\ ,\ which$ shows greater resistance to heat inactivation than other alkaline phosphatase. Presented at high levels in placental trophoblasts, PLAP is recognized as an important marker of differentiation in human malignancies.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




WB result of PLAP antibody (60294-1-Ig, 1:1000) with si-control and si-PLAP transfected HepG2 cells. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




HepG2 cells were subjected to SDS PAGE followed by western blot with 60294-1-1g (PLAP antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60294-1-1g (PLAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




human placenta tissue were subjected to SDS PAGE followed by western blot with 60294-1-lg (PLAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 6024-1-1g (PLAP antibody) at dilution of 1:4000 (under 10x lens). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 60294-1-1g (PLAP antibody) at dilution of 1:4000 (under 40x lens). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 60294-1-lg (PLAP antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60294-1-PBS in a different storage buffer formulation.