## For Research Use Only

## **CREST Monoclonal antibody**

Catalog Number: 60314-1-Ig



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

60314-1-lg

BC034494

Protein G purification

Size:

GeneID (NCBI):

CloneNo.:

1500 µg/ml

26039 **UNIPROT ID:**  3D5D10 Recommended Dilutions:

Source: Mouse

075177

WB 1:1000-1:6000

WB: NIH/3T3 cells, COLO 320 cells, HSC-T6 cells, RAW

Isotype:

Full Name:

lgG1

on chromosome 18-like 1

AG3119

Calculated MW:

synovial sarcoma translocation gene

396 aa. 43 kDa

Observed MW:

50 kDa

**Applications** 

**Tested Applications:** 

Immunogen Catalog Number:

Positive Controls:

WB,ELISA

Species Specificity:

264.7 cells, MCF-7 cells

human, rat, mouse

**Background Information** 

SS18-like 1(SS18L1) is a transcriptional activator that is required for calcium-dependent dendritic growth and branching in cortical neurons. It's also a nuclear protein interacts with CREB-binding protein and expressed in the developing brain. It helps regulate neuronal morphogenesis in calcuim-dependent manner. The N-terminal domain of SS18L1 is required for suppressing transactivation in the basal state, while the C-terminal domain is required for calcium-induced transactivation. It's part of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. This complex also binds to the NR2B promoter, and activity-dependent induction of NR2B expression involves a release of HDAC1 and recruitment of CREBBP. The calculated molecular weight of CREST is about 43 kDa, but the modified of CREST protein is 55 kDa (PMID: 25888396).


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 60314-1-1g (CREST antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.