For Research Use Only

HSP90 Monoclonal antibody, PBS Only

Catalog Number:60318-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number: BC023006

Purification Method: Protein A purification

60318-1-PBS Size:

GeneID (NCBI):

CloneNo.:

1000 µg/ml Source:

3320 **UNIPROT ID:**

P07900

3F11C1

Mouse Isotype:

Full Name:

lgG2a Immunogen Catalog Number:

heat shock protein 90kDa alpha (cytosolic), class A member 1

AG3826

Calculated MW:

853 aa, 90 kDa Observed MW: 85-90 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, CoIP, ELISA

Species Specificity:

human, mouse, rat, pig

Background Information

HSP90, encoded by HSP90AA1, is a constitutively and ubiquitously expressed molecular chaperone that is crucial for $the \, stability \, and \, function \, of \, many \, proteins. \, HSP90 \, provides \, chaperoning \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, many \, of \, them \, activity \, for \, client \, proteins; \, activity \, for$ are members of oncogenic pathways, indicating its implication in tumor malignancy. HSP90 mainly resides in the cytosol, while it can also be released to the extracellular space. Secreted Hsp90 is a C-terminal truncated form. It has been reported that the level of plasma Hsp90 is positively correlated with tumor malignancy in clinical cancer patients, and can be a promising diagnostic marker for tumor malignancy in clinical application.

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only


Selected Validation Data



Co-IP result of anti-HSP90 (IP:60318-1-Ig, 4ug; Detection:13422-1-AP 1:5000) with mouse brain tissue lysate 2240ug. This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60318-1-lg (HSP90 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60318-1-Ig (HSP90 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 60318-1-lg (HSP90 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 60318-1-1g (HSP90 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation



IP result of anti-HSP90 (IP:60318-1-Ig, 5ug; Detection:60318-1-Ig 1:5000) with HepG2 cells lysate 1200 ug. This data was developed using the same antibody clone with 60318-1-PBS in a different storage buffer formulation.