## For Research Use Only

## EEF1B2 Monoclonal antibody

Catalog Number:60329-1-lg Featured Product



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 60329-1-lg
 BC000211

 Size:
 GeneID (NCBI):

 1800 ug/ml
 1933

 Source:
 UNIPROT ID:

 Mouse
 P24534

Isotype: Full Name:
IgG1 eukaryotic translation elongation
factor 1 beta 2

Immunogen Catalog Number: factor 1 beta 2
AG0135 Calculated MW: 25 kDa

Observed MW: 30-34 kDa Purification Method:
Protein G purification

Protein G purification CloneNo.:

3E1C11

Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500 IF/ICC 1:50-1:500

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, RAW 264.7 cells

IHC: human colon tissue,
IF/ICC: HeLa cells,

## **Background Information**

In eukaryotes, the translation elongation factor eEF1A responsible for transporting amino-acylated tRNA to the ribosome forms a higher-order complex, eEF1H, with its guanine-nucleotide-exchange factor eEF1B. eEF1B consists of three subunits: eEF1B alpha, eEF1B beta and eEF1B gamma. The eEF1B2 possess the nucleotide-exchange activity. Although several models on the basis of in vitro experiments have been proposed for the macromolecular organization of the eEF1H complex, these models differ in various aspects. The human eukaryote elongation factor 1 beta 2 (eEF1B2) migrated as a 30-34 kDa protein in SDS-PAGE.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 60329-1-lg (EEF1B2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



WB result of EEF 1B2 antibody (60329-1-1g; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EEF 1B2 transfected HeLa cells.



Immunofluorescent analysis of (-20°C Ethanol ) fixed Hela cells using 60329-1-Ig(EFF1B2 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60329-1-Ig (EEF1B2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).