For Research Use Only

## p63 Monoclonal antibody

Catalog Number:60332-1-lg Featured Product






Basic Information	Catalog Number: 60332-1-lg	GenBank Accession BC039815	n Number:	Purification Method: Protein G purification	
	Size:	GeneID (NCBI):		CloneNo.:	
	633 µg/ml	8626		2B6B2	
	Source: Mouse	UNIPROT ID: Q9H3D4		Recommended Dilutions: WB 1:500-1:2000	
	lsotype: lgG1	Full Name: tumor protein p63		IHC 1:1000-1:4000	
	Immunogen Catalog Number:Calculated MW:AG2791680 aa, 77 kDa				
		Observed MW: 68 kDa			
Applications	Tested Applications: Positive Controls:				
	WB, IHC, ELISA				
	Cited Applications: WB, IHC	Applications: IHC : human tonsillitis tissue.			
	Species Specificity: human				
	Cited Species: human				
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antigen			
Background Information	TP63, also named KET, P63, P73H, P73L and TP73L, belongs to the p53 family. It is a homologue of the tumor suppressor p53 and p73 genes. It is involved in malignancy acquisition and maintenance of cells. Unlike p53, the p63 gene encodes multiple isotypes with remarkably divergent abilities to transactivate p53 reporter genes and induce apoptosis. TP63 acts as a sequence specific DNA binding transcriptional activator or repressor. TP63 has 12 isoforms with MW 40kd(P40), 50kd(P60),63kd(P63) and 73kd(P73L). It is Nuclear stain.				
	Author	Pubmed ID Jou	rnal	Application	
Notable Publications		35238066 And	drologia	IHC	
Notable Publications	Xiqi Ding	55250000 And	-		
Notable Publications			Commun	IHC	
Notable Publications	Li-Yan Li	34272396 Nat	: Commun J Biol Sci	IHC WB	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60332-1-1g (TP63 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of TP63 antibody (60332-1-Ig, 1:1000) with si-control and si-TP63 transfected HepG2 cells.

PC-3 cells were subjected to SDS PAGE followed by western blot with 60332-1-1g (TP63 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

