## For Research Use Only

## Perilipin-2 Monoclonal antibody, PBS Only



Catalog Number: 60340-1-PBS

**Basic Information** 

Catalog Number:

BC005127

**Purification Method:** 

60340-1-PBS

GeneID (NCBI):

Protein A purification CloneNo.:

Size: 1 mg/ml Source:

**UNIPROT ID:** 

1G5D4

Mouse Isotype:

Full Name:

lgG1

adipose differentiation-related

GenBank Accession Number:

protein

Q99541

AG7643

Calculated MW:

48 kDa Observed MW:

47-52 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, Indirect ELISA

Immunogen Catalog Number:

Species Specificity:

human, mouse, rat

## **Background Information**

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT  $family\ which\ is\ responsible\ for\ the\ transportation\ of\ lipids\ and\ the\ formation\ of\ lipid\ droplets.\ ADRP\ is\ localized\ on$ the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


## Selected Validation Data




human adipose tissue were subjected to SDS PAGE followed by western blot with 60340-1-lg (ADRP Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-lg (ADRP Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-1g (ADRP Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human renal cell carcinoma tissue slide using 60340-1-Ig (ADRP/Perilipin 2 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat brown adipose slide using 60340-1-1g (ADRP/Perilipin 2 antibody) at dilution of 1:16000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60340-1-PBS in a different storage buffer formulation.