### For Research Use Only

# BSAP, PAX5 Monoclonal antibody

Catalog Number:60349-1-lg Featured Product

1 Publications

GenBank Accession Number:



**Basic Information** 

Catalog Number: 60349-1-lg

BC156927 GeneID (NCBI): Size: 3100 µg/ml

**UNIPROT ID:** Source: Mouse Q02548 Full Name: Isotype: lgG1 paired box 5 Calculated MW: Immunogen Catalog Number: AG16023 391 aa, 42 kDa

> Observed MW: 42 kDa

**Applications Tested Applications:** 

WB, IHC, FC (Intra), ELISA Cited Applications WB, IHC, IP, IF Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Purification Method:** 

Protein A purification CloneNo.:

2H2D6

Recommended Dilutions: WB 1:500-1:3000 IHC 1:50-1:500

Positive Controls:

WB: Ramos cells, Daudi cells, Raji cells

IHC: human tonsillitis tissue,

## **Background Information**

PAX5, also named B-cell specific activator protein (BSAP), is a member of the highly conserved paired box (PAX)domain family of transcription factors. PAX5 is essential in normal B-cell lymphopoiesis. It is expressed in pro-B, pre-B and mature B lymphocytes but not in plasma cells. PAX5 is a common target in leukemogenesis of B-ALL (Blineage acute lymphoblastic leukemia) along with CDKN2A. Owing to its frequent deletion in B-ALL, PAX5 could be used as one of the molecular markers in diagnosis and monitoring of the disease. This antibody is specific to PAX5. It will not cross react with other PAX members.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yubo Zhao	36064939	Cell Death Dis	WB,IHC,IF,IP

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60349-1-Ig (BSAP,PAX5 Antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60349-1-Ig (BSAP,PAX5 Antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Ramos cells were subjected to SDS PAGE followed by western blot with 60349-1-1g (BSAP,PAX5 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



1X10^6 Ramos cells were intracellularly stained with 0.4 ug Anti-Human BSAP,PAX5 (60349-1-lg, Clone:2H2D6) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).