For Research Use Only

TIM3 Monoclonal antibody, PBS Only

Catalog Number: 60355-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

60355-1-PBS

GeneID (NCBI):

BC020843

Protein A purification

Size: 1mg/ml

GeneID (NCBI 84868 UNIPROT ID: CloneNo.: 4C4G3

Source: Mouse Isotype:

Full Name:

Q8TDQ0

IgG1 hepatitis A virus cellular receptor 2

Immunogen Catalog Number: AG16901

Calculated MW: 301 aa, 33 kDa

Observed MW:

33 kDa, 50-70 kDa

Applications

Tested Applications:

WB, IF-P, ELISA

Species Specificity: human, mouse, rat, pig

Background Information

TIM3, also known as HAVCR2, is a member of the recently discovered T cell Ig and mucin domain-containing molecule superfamily. TIM3 is a negative regulatory molecule that is important for T cell tolerance and has a crucial role in autoimmunity and T cell exhaustion during chronic viral infection (PMID: 23180819). TIM3 is expressed by Thelper type 1 (Th1) cells, macrophage, monocyte, dendritic cells, CD8+ T cell and other lymphocyte subsets. Galectin-9 is a ligand for TIM3. TIM3-galectin-9 pathway negatively regulates T helper type 1 immunity (PMID: 16286920). TIM3 is a 280-aa membrane protein with a calculated molecular weight of 33 kDa, the higher molecular weights between 50 and 70 kDa detected by this monoclonal antibody probably represent glycosylated TIM3 (PMID: 20107545; 17069754; 11725301).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using TIM3 antibody (60355-1-Ig, Clone: 4C4G3) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60355-1-PBS in a different storage buffer formulation.



Daudi cells were subjected to SDS PAGE followed by western blot with 60355-1-1g (TIM3 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60355-1-PBS in a different storage buffer formulation.